

EE595A – Submodular functions, their optimization and applications – Spring 2011

Prof. Jeff Bilmes

University of Washington, Seattle
Department of Electrical Engineering
Spring Quarter, 2011

http://ssli.ee.washington.edu/~bilmes/ee595a_spring_2011/

Lecture 4 - April 8th, 2011

Announcements

- Goal is to have HW1 ready this weekend (so please look out for it).

Matroid

Definition 2.1 (Matroid)

A set system (E, \mathcal{I}) is a **Matroid** if

- (I1') $\emptyset \in \mathcal{I}$
- (I2') $\forall I \in \mathcal{I}, J \subset I \Rightarrow J \in \mathcal{I}$ (down-closed)
- (I3') $\forall I, J \in \mathcal{I}$, with $|I| > |J|$, then there exists $x \in I \setminus J$ such that $J \cup \{x\} \in \mathcal{I}$

Matroids

In fact, we can use the rank of a matroid for its definition.

Theorem 2.2 (Matroid from rank)

Let E be a set and let $r : 2^E \rightarrow \mathbb{Z}_+$ be a function. Then $r(\cdot)$ defines a matroid with r being its rank function if and only if for all $A, B \subseteq E$:

- (R1) $\forall A \subseteq E \quad 0 \leq r(A) \leq |A|$ (non-negative cardinality bounded)
- (R2) $r(A) \leq r(B)$ whenever $A \subseteq B \subseteq E$ (monotone non-decreasing)
- (R3) $r(A \cup B) + r(A \cap B) \leq r(A) + r(B)$ for all $A, B \subseteq E$ (submodular)

- So submodular non-negative integral monotone non-decreasing cardinality bounded is necessary and sufficient to define the matroid.

Matroids

We also saw that it is possible to uniquely define a matroid based on either:

- Independence
- Rank axioms
- Base axioms
- Circuit axioms
- Closure axioms (we didn't see this yesterday, but it is possible)

Matroid, examples

Examples of matroids include

- Matric matroids (characterized by linear independence)
- Graphic matroids (cycle matroid of a graph)
- “free” matroid (all subsets of E)
- k -uniform matroid (all subsets of size at most k)

Partition Matroid

- Let V be our ground set.
- Let $V = V_1 \cup V_2 \cup \dots \cup V_\ell$ be a partition of V into disjoint sets (disjoint union). Define a set of subsets of V as

$$\mathcal{I} = \{X \subseteq V : |X \cap V_i| \leq k_i \text{ for all } i = 1, \dots, \ell\}. \quad (1)$$
 where k_1, \dots, k_ℓ are fixed parameters, $k_i \geq 0$. Then $M = (V, \mathcal{I})$ is a matroid.
- Note that a k -uniform matroid is a trivial example of a partition matroid with $\ell = 1$, $V_1 = V$, and $k_1 = k$.
- We'll show that property (I3') in Def 2.1 holds. If $X, Y \in \mathcal{I}$ with $|Y| > |X|$, then there must be at least one i with $|Y \cap V_i| > |X \cap V_i|$. Therefore, adding one element $e \in V_i \cap (Y \setminus X)$ to X won't break independence.

Partition Matroid

- What is the partition matroid's rank function?

Partition Matroid

- A partition matroids rank function:

$$r(A) = \sum_{i=1}^{\ell} \min(|A \cap V_i|, k_i) \quad (2)$$

which we also immediately see is submodular using properties we spoke about last week.

Partition Matroid

- A partition matroids rank function:

$$r(A) = \sum_{i=1}^{\ell} \min(|A \cap V_i|, k_i) \quad (2)$$

which we also immediately see is submodular using properties we spoke about last week.

- $|A \cap V_i|$ is submodular (even modular) in A

Partition Matroid

- A partition matroids rank function:

$$r(A) = \sum_{i=1}^{\ell} \min(|A \cap V_i|, k_i) \quad (2)$$

which we also immediately see is submodular using properties we spoke about last week.

- $|A \cap V_i|$ is submodular (even modular) in A
- $\min(\text{submodular}(A), k_i)$ is submodular in A since $|A \cap V_i|$ is monotone.

Partition Matroid

- A partition matroids rank function:

$$r(A) = \sum_{i=1}^{\ell} \min(|A \cap V_i|, k_i) \quad (2)$$

which we also immediately see is submodular using properties we spoke about last week.

- $|A \cap V_i|$ is submodular (even modular) in A
- $\min(\text{submodular}(A), k_i)$ is submodular in A since $|A \cap V_i|$ is monotone.
- sums of submodular functions are submodular.

Partition Matroid

- A partition matroids rank function:

$$r(A) = \sum_{i=1}^{\ell} \min(|A \cap V_i|, k_i) \quad (2)$$

which we also immediately see is submodular using properties we spoke about last week.


- $|A \cap V_i|$ is submodular (even modular) in A
- $\min(\text{submodular}(A), k_i)$ is submodular in A since $|A \cap V_i|$ is monotone.
- sums of submodular functions are submodular.
- $r(A)$ is also non-negative integral monotone non-decreasing, so it defines a matroid (the partition matroid).

Partition Matroid, rank as matching

- A partition matroid can be viewed using a bipartite graph.
- Letting V denote the ground set, and V_1, V_2, \dots the partition, the graph is $G = (V, I, E)$ where V is the ground set, I is a set of “indices”, and E is the set of edges.
- $I = (I_1, I_2, \dots, I_\ell)$ is a set of $k = \sum_{i=1}^{\ell} k_i$ nodes, grouped into ℓ clusters, where there are k_i nodes in the i^{th} group I_i .
- $(v, i) \in E(G)$ iff $v \in V_j$ and $i \in I_j$.


Partition Matroid, rank as matching

- Example where $\ell = 5$,
 $(k_1, k_2, k_3, k_4, k_5) =$
 $(2, 2, 1, 1, 3)$.



Partition Matroid, rank as matching


- Example where $\ell = 5$,
 $(k_1, k_2, k_3, k_4, k_5) =$
 $(2, 2, 1, 1, 3)$.



- Recall, $\Gamma : 2^V \rightarrow \mathbb{R}$ as the neighbor function in a bipartite graph, the neighbors of X is defined as $\Gamma(X) = \{v \in V(G) \setminus X : E(X, \{v\}) \neq \emptyset\}$, and how $|\Gamma(X)|$ is submodular.

Partition Matroid, rank as matching


- Example where $\ell = 5$,
 $(k_1, k_2, k_3, k_4, k_5) =$
 $(2, 2, 1, 1, 3)$.



- Recall, $\Gamma : 2^V \rightarrow \mathbb{R}$ as the neighbor function in a bipartite graph, the neighbors of X is defined as $\Gamma(X) = \{v \in V(G) \setminus X : E(X, \{v\}) \neq \emptyset\}$, and how $|\Gamma(X)|$ is submodular.
- Here, for $X \subseteq V$, we have $\Gamma(X) = \{i \in I : (v, i) \in E(G) \text{ and } v \in X\}$.

Partition Matroid, rank as matching

- Example where $\ell = 5$,
 $(k_1, k_2, k_3, k_4, k_5) =$
 $(2, 2, 1, 1, 3)$.



- Recall, $\Gamma : 2^V \rightarrow \mathbb{R}$ as the neighbor function in a bipartite graph, the neighbors of X is defined as $\Gamma(X) = \{v \in V(G) \setminus X : E(X, \{v\}) \neq \emptyset\}$, and how $|\Gamma(X)|$ is submodular.
- Here, for $X \subseteq V$, we have $\Gamma(X) = \{i \in I : (v, i) \in E(G) \text{ and } v \in X\}$.
- For such a constructed bi-partite graph, the rank function of a partition matroid is

$$r(X) = \sum_{i=1}^{\ell} \min(|X \cap V_i|, k_i) =$$
maximum matching involving X .

System of Representatives

- Let (V, \mathcal{V}) be a set system (i.e., $\mathcal{V} = (V_k : k \in I)$ where $V_i \subseteq V$ for all i).

System of Representatives

- Let (V, \mathcal{V}) be a set system (i.e., $\mathcal{V} = (V_k : k \in I)$ where $V_i \subseteq V$ for all i).
- A family $(v_i : i \in I)$ for index set I is said to be a **system of representatives** of \mathcal{V} if \exists a bijection $\pi : I \rightarrow I$ such that $v_i \in V_{\pi(i)}$. *v_i is the representative of set $\pi(i)$, meaning the i^{th} representative is meant to represent set $V_{\pi(i)}$. Consider the house of representatives, $v_i = \text{"John Smith"}$, while $i = \text{King County}$.*

System of Representatives


- Let (V, \mathcal{V}) be a set system (i.e., $\mathcal{V} = (V_k : k \in I)$ where $V_i \subseteq V$ for all i).
- A family $(v_i : i \in I)$ for index set I is said to be a **system of representatives** of \mathcal{V} if \exists a bijection $\pi : I \rightarrow I$ such that $v_i \in V_{\pi(i)}$.
- In a system of representatives, there is no requirement for the representatives to be distinct. I.e., we could have $v_1 \in T$, where v_1 represents both V_1 and V_2 .

System of Representatives

- Let (V, \mathcal{V}) be a set system (i.e., $\mathcal{V} = (V_k : k \in I)$ where $V_i \subseteq V$ for all i).
- A family $(v_i : i \in I)$ for index set I is said to be a **system of representatives** of \mathcal{V} if \exists a bijection $\pi : I \rightarrow I$ such that $v_i \in V_{\pi(i)}$.
- In a system of representatives, there is no requirement for the representatives to be distinct. I.e., we could have $v_1 \in T$, where v_1 represents both V_1 and V_2 .
- We can view this as a bipartite graph.


System of Representatives

- We can view this as a bipartite graph. The groups of V are marked by color tags on the left, and also via right neighbors in the graph.
- Here, $\ell = 6$, and $\mathcal{V} = (V_1, V_2, \dots, V_6)$
 $= (\{e, f, h\}, \{d, e, g\}, \{b, c, e, h\}, \{a, b, h\}, \{a\}, \{a\})$.



System of Representatives


- We can view this as a bipartite graph. The groups of V are marked by color tags on the left, and also via right neighbors in the graph.
- Here, $\ell = 6$, and $\mathcal{V} = (V_1, V_2, \dots, V_6)$
 $= (\{e, f, h\}, \{d, e, g\}, \{b, c, e, h\}, \{a, b, h\}, \{a\}, \{a\})$.



- A system of representatives would make sure that there is a representative for each color group. For example,

System of Representatives


- We can view this as a bipartite graph. The groups of V are marked by color tags on the left, and also via right neighbors in the graph.
- Here, $\ell = 6$, and $\mathcal{V} = (V_1, V_2, \dots, V_6)$
 $= (\{e, f, h\}, \{d, e, g\}, \{b, c, e, h\}, \{a, b, h\}, \{a\}, \{a\})$.



- A system of representatives would make sure that there is a representative for each color group. For example,
- The representatives are shown as colors on the left.

System of Representatives

- We can view this as a bipartite graph. The groups of V are marked by color tags on the left, and also via right neighbors in the graph.
- Here, $\ell = 6$, and $\mathcal{V} = (V_1, V_2, \dots, V_6)$
 $= (\{e, f, h\}, \{d, e, g\}, \{b, c, e, h\}, \{a, b, h\}, \{a\}, \{a\})$.



- A system of representatives would make sure that there is a representative for each color group. For example,
- The representatives are shown as colors on the left.
- Here, the set of representatives is not distinct. In fact, due to the red and pink group, a distinct group of representatives is impossible (since there is only one common choice to represent both color groups).

System of Distinct Representatives

- Let (V, \mathcal{V}) be a set system (i.e., $\mathcal{V} = (V_k : k \in I)$ where $V_i \subseteq V$ for all i).

System of Distinct Representatives

- Let (V, \mathcal{V}) be a set system (i.e., $\mathcal{V} = (V_k : k \in I)$ where $V_i \subseteq V$ for all i).
- A family $(v_i : i \in I)$ for index set I is said to be a **system of distinct representatives** of \mathcal{V} if \exists a bijection $\pi : I \leftrightarrow I$ such that $v_i \in V_{\pi(i)}$ and $v_i \neq v_j$ for all $i \neq j$.

System of Distinct Representatives

- Let (V, \mathcal{V}) be a set system (i.e., $\mathcal{V} = (V_k : k \in I)$ where $V_i \subseteq V$ for all i).
- A family $(v_i : i \in I)$ for index set I is said to be a **system of distinct representatives** of \mathcal{V} if \exists a bijection $\pi : I \leftrightarrow I$ such that $v_i \in V_{\pi(i)}$ and $v_i \neq v_j$ for all $i \neq j$.
- In a system of distinct representatives, there **is** a requirement for the representatives to be distinct.

System of Distinct Representatives

- Let (V, \mathcal{V}) be a set system (i.e., $\mathcal{V} = (V_k : i \in I)$ where $V_i \subseteq V$ for all i).
- A family $(v_i : i \in I)$ for index set I is said to be a **system of distinct representatives** of \mathcal{V} if \exists a bijection $\pi : I \leftrightarrow I$ such that $v_i \in V_{\pi(i)}$ and $v_i \neq v_j$ for all $i \neq j$.
- In a system of distinct representatives, there **is** a requirement for the representatives to be distinct.

Definition 4.1 (transversal)

Given a set system (V, \mathcal{V}) , a set $T \subseteq V$ is a **transversal** of \mathcal{V} if there is a bijection $\pi : T \leftrightarrow I$ such that

$$x \in V_{\pi(x)} \text{ for all } x \in T \quad (3)$$

System of Distinct Representatives

- Let (V, \mathcal{V}) be a set system (i.e., $\mathcal{V} = (V_k : i \in I)$ where $V_i \subseteq V$ for all i).
- A family $(v_i : i \in I)$ for index set I is said to be a **system of distinct representatives** of \mathcal{V} if \exists a bijection $\pi : I \leftrightarrow I$ such that $v_i \in V_{\pi(i)}$ and $v_i \neq v_j$ for all $i \neq j$.
- In a system of distinct representatives, there **is** a requirement for the representatives to be distinct.

Definition 4.1 (transversal)

Given a set system (V, \mathcal{V}) , a set $T \subseteq V$ is a **transversal** of \mathcal{V} if there is a bijection $\pi : T \leftrightarrow I$ such that

$$x \in V_{\pi(x)} \text{ for all } x \in T \quad (3)$$

- Note that due to it being a bijection, all of I and T are “covered” (so this makes things distinct).

Transversal

- A set $X \subseteq V$ is a **partial transversal** if X is a transversal of some subfamily $\mathcal{V}' = (V_k : i \in I')$ where $I' \subseteq I$.

Transversal

- A set $X \subseteq V$ is a **partial transversal** if X is a transversal of some subfamily $\mathcal{V}' = (V_k : i \in I')$ where $I' \subseteq I$.
- Therefore, for any transversal T , any subset $T' \subseteq T$ is a partial transversal (down closed).

When Transversals?

- As we saw, a transversal might not always exist. In fact, we have the following theorems.

When Transversals?

- As we saw, a transversal might not always exist. In fact, we have the following theorems.
- Given a set system (V, \mathcal{V}) with $\mathcal{V} = (V_i : i \in I)$, for any $J \subseteq I$, let
$$V(J) = \cup_{j \in J} V_j \tag{4}$$

When Transversals?

- As we saw, a transversal might not always exist. In fact, we have the following theorems.
- Given a set system (V, \mathcal{V}) with $\mathcal{V} = (V_i : i \in I)$, for any $J \subseteq I$, let

$$V(J) = \cup_{j \in J} V_j \tag{4}$$
- We have

Theorem 5.1 (Hall's theorem)


Given a set system (V, \mathcal{V}) , the family of subsets $\mathcal{V} = (V_i : i \in I)$ has a transversal iff for all $J \subseteq I$

$$|V(J)| \geq |J| \tag{5}$$

When Transversals?

- As we saw, a transversal might not always exist. In fact, we have the following theorems.
- Given a set system (V, \mathcal{V}) with $\mathcal{V} = (V_i : i \in I)$, for any $J \subseteq I$, let

$$V(J) = \cup_{j \in J} V_j \quad (4)$$
- Hall's theorem as a bipartite graph.



When Transversals?

- As we saw, a transversal might not always exist. In fact, we have the following theorems.
- Given a set system (V, \mathcal{V}) with $\mathcal{V} = (V_i : i \in I)$, for any $J \subseteq I$, let

$$V(J) = \cup_{j \in J} V_j \quad (4)$$
- Moreover, we have

Theorem 5.2 (Rado's theorem)

If $M = (V, r)$ is a matroid on V with rank function r , then the family of subsets $(V_i : i \in I)$ of V has a transversal which is independent in M iff for all $J \subseteq I$

$$r(V(J)) \geq |J| \quad (6)$$

When Transversals?

- As we saw, a transversal might not always exist. In fact, we have the following theorems.
- Given a set system (V, \mathcal{V}) with $\mathcal{V} = (V_i : i \in I)$, for any $J \subseteq I$, let

$$V(J) = \cup_{j \in J} V_j \quad (4)$$
- Moreover, we have

Theorem 5.2 (Rado's theorem)

If $M = (V, r)$ is a matroid on V with rank function r , then the family of subsets $(V_i : i \in I)$ of V has a transversal which is independent in M iff for all $J \subseteq I$

$$r(V(J)) \geq |J| \quad (6)$$

- Note, a transversal T independent in M means that $r(T') = |T'|$ for all $T' \subseteq T$.

When Transversals?

Theorem 5.3

If $\mathcal{V} = (V_i : i \in I)$ is a finite family of non-empty subsets of V , and $f : 2^V \rightarrow \mathbb{Z}_+$ is a non-negative, integral, monotone non-decreasing, and submodular function, then \mathcal{V} has a system of representatives $(v_i : i \in I)$ such that

$$f(\cup_{i \in J} \{v_i\}) \geq |J| \text{ for all } J \subseteq I \quad (7)$$

if and only if

$$f(V(J)) \geq |J| \text{ for all } J \subseteq I \quad (8)$$

When Transversals?

Theorem 5.3

If $\mathcal{V} = (V_i : I \in I)$ is a finite family of non-empty subsets of V , and $f : 2^V \rightarrow \mathbb{Z}_+$ is a non-negative, integral, monotone non-decreasing, and submodular function, then \mathcal{V} has a system of representatives $(v_i : i \in I)$ such that

$$f(\cup_{i \in J} \{v_i\}) \geq |J| \text{ for all } J \subseteq I \quad (7)$$

if and only if

$$f(V(J)) \geq |J| \text{ for all } J \subseteq I \quad (8)$$

- Given Theorem 5.3, we immediately get Theorem 5.1 by taking $f(S) = |S|$ for $S \subseteq V$. *In which case, Eq. 7 requires the system of representatives to be distinct.*

When Transversals?

Theorem 5.3

If $\mathcal{V} = (V_i : I \in I)$ is a finite family of non-empty subsets of V , and $f : 2^V \rightarrow \mathbb{Z}_+$ is a non-negative, integral, monotone non-decreasing, and submodular function, then \mathcal{V} has a system of representatives $(v_i : i \in I)$ such that

$$f(\cup_{i \in J} \{v_i\}) \geq |J| \text{ for all } J \subseteq I \quad (7)$$

if and only if

$$f(V(J)) \geq |J| \text{ for all } J \subseteq I \quad (8)$$

- Given Theorem 5.3, we immediately get Theorem 5.1 by taking $f(S) = |S|$ for $S \subseteq V$.
- We get Theorem 5.2 by taking $f(S) = r(S)$ for $S \subseteq V$, the rank function of the matroid. *where, Eq. 7 insists the system of representatives is independent in M .*

When Transversals?

first part proof of Theorem 5.3.

- Suppose Eq. 7 is true. Then since f is monotone, and since $V(J) \supseteq \cup_{i \in J} \{v_i\}$ when $(v_i : i \in I)$ is a system of representatives, then Eq. 8 immediately follows.

...

When Transversals?

Lemma 5.4

Suppose Eq. 8 is true for \mathcal{V} , and there exists an i such that $|V_i| > 2$. W.l.o.g. let $i = 1$. Then there exists $f \in V_1$ such that the family of subsets $(V_1 \setminus \{f\}, V_2, \dots, V_n)$ also satisfies Eq 8.

Proof.

- When Eq. 8 holds, this means that for any subsets $J_1, J_2 \subseteq \{2, \dots, n\}$, we have that

$$f(V_1 \cup V(J_1)) \geq |J_1| + 1 \tag{9}$$

$$f(V_1 \cup V(J_2)) \geq |J_2| + 1 \tag{10}$$

...

When Transversals?

Lemma 5.4

Suppose Eq. 8 is true for \mathcal{V} , and there exists an i such that $|V_i| > 2$. W.l.o.g. let $i = 1$. Then there exists $f \in V_1$ such that the family of subsets $(V_1 \setminus \{f\}, V_2, \dots, V_n)$ also satisfies Eq 8.

Proof.

- Suppose, to the contrary, this is false. Then we may take f_1, f_2 as two distinct elements in V_1 ,

...

When Transversals?

Lemma 5.4

Suppose Eq. 8 is true for \mathcal{V} , and there exists an i such that $|V_i| > 2$. W.l.o.g. let $i = 1$. Then there exists $f \in V_1$ such that the family of subsets $(V_1 \setminus \{f\}, V_2, \dots, V_n)$ also satisfies Eq 8.

Proof.

- Suppose, to the contrary, this is false. Then we may take f_1, f_2 as two distinct elements in V_1 ,
- and there must exist subsets J_1, J_2 of $\{2, \dots, n\}$ such that

$$f((V_1 \setminus \{f_1\}) \cup V(J_1)) < |J_1| + 1, \quad (11)$$

$$f((V_1 \setminus \{f_2\}) \cup V(J_2)) < |J_2| + 1, \quad (12)$$

(note that either one or both of J_1, J_2 could be empty).

...

When Transversals?

Lemma 5.4

Suppose Eq. 8 is true for \mathcal{V} , and there exists an i such that $|V_i| > 2$. W.l.o.g. let $i = 1$. Then there exists $f \in V_1$ such that the family of subsets $(V_1 \setminus \{f\}, V_2, \dots, V_n)$ also satisfies Eq 8.

Proof.

- Taking $X = (V_1 \setminus \{f_1\}) \cup V(J_1)$ and $Y = (V_1 \setminus \{f_2\}) \cup V(J_2)$, we see that:

$$X \cup Y = V_1 \cup V(J_1 \cup J_2) \quad (13)$$

and

$$X \cap Y \supseteq V(J_1 \cap J_2) \quad (14)$$

...

When Transversals?

Lemma 5.4

Suppose Eq. 8 is true for \mathcal{V} , and there exists an i such that $|V_i| > 2$. W.l.o.g. let $i = 1$. Then there exists $f \in V_1$ such that the family of subsets $(V_1 \setminus \{f\}, V_2, \dots, V_n)$ also satisfies Eq 8.

Proof.

- since f is submodular and monotone non-decreasing, we get

$$|J_1| + |J_2| \geq f(V_1 \cup V(J_1 \cup J_2)) + f(V(J_1 \cap J_2)) \quad (15)$$

...

When Transversals?

Lemma 5.4

Suppose Eq. 8 is true for \mathcal{V} , and there exists an i such that $|V_i| > 2$. W.l.o.g. let $i = 1$. Then there exists $f \in V_1$ such that the family of subsets $(V_1 \setminus \{f\}, V_2, \dots, V_n)$ also satisfies Eq 8.

Proof.

- since f is submodular and monotone non-decreasing, we get

$$|J_1| + |J_2| \geq f(V_1 \cup V(J_1 \cup J_2)) + f(V(J_1 \cap J_2)) \quad (15)$$

- Since \mathcal{V} satisfies Eq. 8, and $1 \notin J_1 \cup J_2$, this gives

$$|J_1| + |J_2| \geq |J_1 \cup J_2| + 1 + |J_1 \cap J_2| \quad (16)$$

which is a contradiction.



When Transversals?

converse proof of Theorem 5.3.

- Conversely, suppose Eq. 8 is true.



When Transversals?

converse proof of Theorem 5.3.

- Conversely, suppose Eq. 8 is true.
- If each V_i is a singleton set, then the result follows immediately.



When Transversals?

converse proof of Theorem 5.3.

- Conversely, suppose Eq. 8 is true.
- If each V_i is a singleton set, then the result follows immediately.
- W.l.o.g. , let $|V_1| \geq 2$, then by Lemma 5.4, the family of subsets $(V_1 \setminus \{f\}, V_2, \dots, V_n)$ also satisfies Eq 8.



When Transversals?

converse proof of Theorem 5.3.

- Conversely, suppose Eq. 8 is true.
- If each V_i is a singleton set, then the result follows immediately.
- W.l.o.g. , let $|V_1| \geq 2$, then by Lemma 5.4, the family of subsets $(V_1 \setminus \{f\}, V_2, \dots, V_n)$ also satisfies Eq 8.
- We can continue to reduce the family, deleting elements from V_i for some i until we arrive at a family of singleton sets.



When Transversals?

converse proof of Theorem 5.3.

- Conversely, suppose Eq. 8 is true.
- If each V_i is a singleton set, then the result follows immediately.
- W.l.o.g. , let $|V_1| \geq 2$, then by Lemma 5.4, the family of subsets $(V_1 \setminus \{f\}, V_2, \dots, V_n)$ also satisfies Eq 8.
- We can continue to reduce the family, deleting elements from V_i for some i until we arrive at a family of singleton sets.
- This family will be the required system of representatives.



When Transversals?

converse proof of Theorem 5.3.

- Conversely, suppose Eq. 8 is true.
- If each V_i is a singleton set, then the result follows immediately.
- W.l.o.g. , let $|V_1| \geq 2$, then by Lemma 5.4, the family of subsets $(V_1 \setminus \{f\}, V_2, \dots, V_n)$ also satisfies Eq 8.
- We can continue to reduce the family, deleting elements from V_i for some i until we arrive at a family of singleton sets.
- This family will be the required system of representatives.



This theorem can be used to produce a variety of other results quite easily, and shows how submodularity is the key ingredient in its truth.

Transversal Matroid

Transversals, themselves, define a matroid.

Theorem 6.1

If \mathcal{V} is a family of finite subsets of a ground set V , then the collection of partial transversals of \mathcal{V} is the set of independent sets of a matroid $M = (V, \mathcal{V})$ on V .

Transversal Matroid

Transversals, themselves, define a matroid.

Theorem 6.1

If \mathcal{V} is a family of finite subsets of a ground set V , then the collection of partial transversals of \mathcal{V} is the set of independent sets of a matroid $M = (V, \mathcal{V})$ on V .

- This means that the transversals of \mathcal{V} are the bases of matroid M . Therefore, all maximal partial transversals of \mathcal{V} have the same cardinality!

Transversals and Matchings

- Transversals correspond exactly to matchings in bipartite graphs (as we've already strongly hinted at).

Transversals and Matchings

- Transversals correspond exactly to matchings in bipartite graphs (as we've already strongly hinted at).
- Given a set system (V, \mathcal{V}) , with $\mathcal{V} = (V_i : i \in I)$, we can define a bipartite graph $G = (V, I, E)$ associated with \mathcal{V} that has edge set $\{(v, i) : v \in V, i \in I, v \in V_i\}$.

Transversals and Matchings

- Transversals correspond exactly to matchings in bipartite graphs (as we've already strongly hinted at).
- Given a set system (V, \mathcal{V}) , with $\mathcal{V} = (V_i : i \in I)$, we can define a bipartite graph $G = (V, I, E)$ associated with \mathcal{V} that has edge set $\{(v, i) : v \in V, i \in I, v \in I_i\}$.
- A **matching** in this graph is a set of edges no two of which have a common endpoint.

Transversals and Matchings

- Transversals correspond exactly to matchings in bipartite graphs (as we've already strongly hinted at).
- Given a set system (V, \mathcal{V}) , with $\mathcal{V} = (V_i : i \in I)$, we can define a bipartite graph $G = (V, I, E)$ associated with \mathcal{V} that has edge set $\{(v, i) : v \in V, i \in I, v \in I_i\}$.
- A **matching** in this graph is a set of edges no two of which that have a common endpoint.
- In fact, we easily have

Lemma 6.2

A subset $T \subseteq V$ is a partial transversal of \mathcal{V} iff there is a matching in (V, I, E) in which every edge has one endpoint in T .

We say that T is matched into I .

Morphing Partition Matroid Rank

- Recall the partition matroid rank function. Note also that $k_i = |I_i|$ in the bipartite graph representation.

Morphing Partition Matroid Rank

- Recall the partition matroid rank function. Note also that $k_i = |I_i|$ in the bipartite graph representation.
- We change the form of this in the subsequent equations.

$$r(A) = \sum_{i=1}^{\ell} \min(|A \cap V_i|, k_i) \quad (17)$$

Morphing Partition Matroid Rank

- Recall the partition matroid rank function. Note also that $k_i = |I_i|$ in the bipartite graph representation.
- We change the form of this in the subsequent equations.

$$r(A) = \sum_{i=1}^{\ell} \min(|A \cap V_i|, k_i) \quad (17)$$

$$= \sum_{i=1}^{\ell} \min(|A \cap V(I_i)|, |I_i|) \quad (18)$$

Morphing Partition Matroid Rank

- Recall the partition matroid rank function. Note also that $k_i = |I_i|$ in the bipartite graph representation.
- We change the form of this in the subsequent equations.

$$r(A) = \sum_{i=1}^{\ell} \min(|A \cap V_i|, k_i) \quad (17)$$

$$= \sum_{i=1}^{\ell} \min(|A \cap V(I_i)|, |I_i|) \quad (18)$$

$$= \sum_{i=1}^{\ell} \min_{J_i \subseteq I_i} \left(\left\{ \begin{array}{ll} |A \cap V(I_i)| & \text{if } J_i \neq \emptyset \\ 0 & \text{if } J_i = \emptyset \end{array} \right\} + |I_i \setminus J_i| \right) \quad (19)$$

Morphing Partition Matroid Rank

- Recall the partition matroid rank function. Note also that $k_i = |I_i|$ in the bipartite graph representation.
- We change the form of this in the subsequent equations.

$$r(A) = \sum_{i=1}^{\ell} \min(|A \cap V_i|, k_i) \quad (17)$$

$$= \sum_{i=1}^{\ell} \min(|A \cap V(I_i)|, |I_i|) \quad (18)$$

$$= \sum_{i=1}^{\ell} \min_{J_i \subseteq I_i} \left(\begin{cases} |A \cap V(I_i)| & \text{if } J_i \neq \emptyset \\ 0 & \text{if } J_i = \emptyset \end{cases} + |I_i \setminus J_i| \right) \quad (19)$$

$$= \sum_{i=1}^{\ell} \min_{J_i \subseteq I_i} (|V(J_i) \cap A| + |I_i \setminus J_i|) \quad (20)$$

... Morphing Partition Matroid Rank

- Continuing,

$$r(A) = \sum_{i=1}^{\ell} \min_{J_i \subseteq I_i} (|V(J_i) \cap V(I_i) \cap A| - |I_i \cap J_i| + |I_i|) \quad (21)$$

... Morphing Partition Matroid Rank

- Continuing,

$$r(A) = \sum_{i=1}^{\ell} \min_{J_i \subseteq I_i} (|V(J_i) \cap V(I_i) \cap A| - |I_i \cap J_i| + |I_i|) \quad (21)$$

$$= \min_{J \subseteq I} \left(\sum_{i=1}^{|I|} |V(J) \cap V(I_i) \cap A| - |I_i \cap J| + |I_i| \right) \quad (22)$$

... Morphing Partition Matroid Rank

- Continuing,

$$r(A) = \sum_{i=1}^{\ell} \min_{J_i \subseteq I_i} (|V(J_i) \cap V(I_i) \cap A| - |I_i \cap J_i| + |I_i|) \quad (21)$$

$$= \min_{J \subseteq I} \left(\sum_{i=1}^{|I|} |V(J) \cap V(I_i) \cap A| - |I_i \cap J| + |I_i| \right) \quad (22)$$

$$= \min_{J \subseteq I} (|V(J) \cap V(I) \cap A| - |J| + |I|) \quad (23)$$

... Morphing Partition Matroid Rank

- Continuing,

$$r(A) = \sum_{i=1}^{\ell} \min_{J_i \subseteq I_i} (|V(J_i) \cap V(I_i) \cap A| - |I_i \cap J_i| + |I_i|) \quad (21)$$

$$= \min_{J \subseteq I} \left(\sum_{i=1}^{|I|} |V(J) \cap V(I_i) \cap A| - |I_i \cap J| + |I_i| \right) \quad (22)$$

$$= \min_{J \subseteq I} (|V(J) \cap V(I) \cap A| - |J| + |I|) \quad (23)$$

$$= \min_{J \subseteq I} (|V(J) \cap A| - |J| + |I|) \quad (24)$$

... Morphing Partition Matroid Rank

- Continuing,

$$r(A) = \sum_{i=1}^{\ell} \min_{J_i \subseteq I_i} (|V(J_i) \cap V(I_i) \cap A| - |I_i \cap J_i| + |I_i|) \quad (21)$$

$$= \min_{J \subseteq I} \left(\sum_{i=1}^{|I|} |V(J) \cap V(I_i) \cap A| - |I_i \cap J| + |I_i| \right) \quad (22)$$

$$= \min_{J \subseteq I} (|V(J) \cap V(I) \cap A| - |J| + |I|) \quad (23)$$

$$= \min_{J \subseteq I} (|V(J) \cap A| - |J| + |I|) \quad (24)$$

- In fact, this bottom (more general) expression is the expression for the rank of a transversal matroid.

Partial Transversals Are Matroids

In fact, we have

Theorem 6.3

Let (V, \mathcal{V}) where $\mathcal{V} = (V_1, V_2, \dots, V_\ell)$ be a subset system. Let $I = \{1, \dots, \ell\}$. Let \mathcal{I} be the set of partial transversals of \mathcal{V} . Then (V, \mathcal{I}) is a matroid.

Proof.



Partial Transversals Are Matroids

In fact, we have

Theorem 6.3

Let (V, \mathcal{V}) where $\mathcal{V} = (V_1, V_2, \dots, V_\ell)$ be a subset system. Let $I = \{1, \dots, \ell\}$. Let \mathcal{I} be the set of partial transversals of \mathcal{V} . Then (V, \mathcal{I}) is a matroid.

Proof.

- We note that $\emptyset \in \mathcal{I}$ since the empty set is a transversal of the empty subfamily of \mathcal{V} , thus (I1') holds.



Partial Transversals Are Matroids

In fact, we have

Theorem 6.3

Let (V, \mathcal{V}) where $\mathcal{V} = (V_1, V_2, \dots, V_\ell)$ be a subset system. Let $I = \{1, \dots, \ell\}$. Let \mathcal{I} be the set of partial transversals of \mathcal{V} . Then (V, \mathcal{I}) is a matroid.

Proof.

- We note that $\emptyset \in \mathcal{I}$ since the empty set is a transversal of the empty subfamily of \mathcal{V} , thus (I1') holds.
- We already saw that if T is a partial transversal of \mathcal{V} , and if $T' \subseteq T$, then T' is also a partial transversal. So (I2') holds.



Partial Transversals Are Matroids

In fact, we have

Theorem 6.3

Let (V, \mathcal{V}) where $\mathcal{V} = (V_1, V_2, \dots, V_\ell)$ be a subset system. Let $I = \{1, \dots, \ell\}$. Let \mathcal{I} be the set of partial transversals of \mathcal{V} . Then (V, \mathcal{I}) is a matroid.

Proof.

- We note that $\emptyset \in \mathcal{I}$ since the empty set is a transversal of the empty subfamily of \mathcal{V} , thus (I1') holds.
- We already saw that if T is a partial transversal of \mathcal{V} , and if $T' \subseteq T$, then T' is also a partial transversal. So (I2') holds.
- Suppose that T_1 and T_2 are partial transversals of \mathcal{V} such that $|T_1| < |T_2|$. **Exercise: show that (I3') holds.**



Transversal Matroid Rank

- Transversal matroid has rank

$$r(A) = \min_{J \subseteq I} (|V(J) \cap A| - |J| + |I|) \quad (25)$$

Transversal Matroid Rank

- Transversal matroid has rank

$$r(A) = \min_{J \subseteq I} (|V(J) \cap A| - |J| + |I|) \quad (25)$$

- Therefore, this function is submodular.

Transversal Matroid Rank

- Transversal matroid has rank

$$r(A) = \min_{J \subseteq I} (|V(J) \cap A| - |J| + |I|) \quad (25)$$

- Therefore, this function is submodular.
- Note that it is a minimum over a set of modular functions. Is this true in general? **Exercise:**

Matroid and the greedy algorithm

- Let \mathcal{I} be a set of subsets of E that is down-closed. Consider a modular weight function $w : E \rightarrow \mathbb{R}$, and we want to find the $A \in \mathcal{I}$ that maximizes $w(A)$.
- Greedy algorithm: Set $A = \emptyset$, and repeatedly choose $y \in E \setminus A$ such that $A \cup \{y\} \in \mathcal{I}$ with $w(y)$ as large as possible, stopping when no such y exists.

Theorem 7.1

Let \mathcal{I} be a non-empty collection of subsets of a set E , down-closed. Then the pair (E, \mathcal{I}) is a matroid if and only if for each weight function $w \in \mathcal{R}^E$, the greedy algorithm leads to a set $I \in \mathcal{I}$ of maximum weight $w(I)$.

Scratch Paper

Scratch Paper

Scratch Paper

Sources for Today's Lecture

Korte, Vygen-2005, Vondrak-2010, Schrijver-2003, Oxley-1992, Welsh-1976, Goemans-2010.